

0535

Lemna  
duckweed  
macroalgae  
microalgae

REVIEW

Literature Review on Duckweed Toxicity Testing

WUNCHENG WANG

Water Quality Section, Illinois State Water Survey, Box 697, Peoria, Illinois 61652

Received October 20, 1989

Duckweed commonly refers to a group of floating, flowering plants of the family Lemnaceae. Duckweed plants are fast growing and widely distributed. They are easy to culture and to test. Some reports suggest that duckweed plants are tolerant to environmental toxicity. Other studies, however, indicate that duckweed plants are as sensitive to toxicity as other aquatic species. Duckweed plants are especially suitable for use in complex effluent bioassays, and for testing herbicide pollution in the aquatic environment, lake and river pollution, sediment toxicity, and the like. Duckweed and algae represent different levels of complexity in the plant kingdom. They complement each other as phytotoxicity test organisms, instead of mutually excluding each other. Many duckweed species have been studied, primarily of the *Lemna* and *Spirodela* genera. *Lemna minor* and *L. gibba* have been recommended as standard test species. Differences in duckweed test methodology occur with regard to test types, test vessels, control tests, nutrient media, end points, and applications. © 1990 Academic Press, Inc.

INTRODUCTION

Three base-set tests are currently used for freshwater environmental monitoring, effluent toxicity testing, toxicity assessment of a product, etc. They are tests of *Pimephales promelas*, *Daphnia magna* (or *Daphnia pulex*), and *Selenastrum capricornutum* (Peltier and Weber, 1985; Horning and Weber, 1985). These tests are generally applicable to important environmental laws such as the National Pollutant Discharge Elimination System regulations, the Toxic Substance Control Act, and the like. They are used routinely by federal and state regulatory agencies, industries, and consulting laboratories.

Recently the duckweed toxicity test has received much attention. Duckweed is an aquatic plant and is relevant to many aquatic environments, including lakes, streams, effluent, rain, and sediment. Additionally, duckweed is a vascular, flowering plant which provides additional information unlikely to be obtained by the three base-set tests.

The objective of this article is a critical review on the duckweed toxicity test. It is intended to be methodology oriented, rather than an exhaustive literature citation. Other related review articles on duckweed are also available (Hillman, 1961; Hillman and Culley, 1978; Wang, submitted for publication).

FEATURES OF DUCKWEED PLANTS

The term "duckweed" commonly refers to a group of aquatic vascular angiosperms of the family Lemnaceae. Duckweed plants are divided into four genera, *Spirodela*, *Lemna*, *Wolffiella*, and *Wolffia*. There are approximately 40 spe-

cies worldwide, half of which are found in the United States. Duckweed plants are widely distributed in the world from the tropical to the temperate zones, from freshwater to brackish estuaries, and throughout a wide range of trophic conditions (Hillman and Culley, 1978). Duckweed plants are common in the aquatic environment, especially in quiescent water bodies.

Duckweed plants are composed of two parts, frond and root. The plants are colonial and form aggregates of two or more fronds in a colony. *Lemna minor* has a single rootlet, while *Spirodela* has several rootlets (Correll and Correll, 1972).

Duckweed plants are small. *L. minor* is 2–4 mm across, while *Spirodela* thallus is 3–10 mm long (Correll and Correll, 1972). The plant size is small so that a large laboratory space is not required for culturing or testing. Yet the size is sufficiently large to be visually observed, allowing nondestructive, repeated observations.

Duckweed plants are extremely fast growing (Hillman and Culley, 1978). The plants form a thick mat, frequently dominated by a single species, in a lake or a pond. Fish kills have been reported as a result of duckweed and algal domination (Lewis and Bender, 1961). In an 18-month study, the doubling time for *L. minor* fronds ranged from 1.3 to 2.8 days, as shown in Fig. 1 (Wang, 1987a). The mean value and standard deviation were 1.9 and 0.36 days, respectively. This series of experiments were performed using a nutrient solution containing a double-strength (2×) algal nutrient solution (American Public Health Association *et al.*, 1985). Frick (1985) reported that the frond doubling time for *L. minor* was about 1.4 days. In comparison, Hughes *et al.* (1988) reported the doubling time for *L. gibba* to be 0.7 days. On the basis of results reported by Nasu *et al.* (1984), the doubling time for *L. paucicostata* was calculated to be 0.35 days. Duckweed cultured in the laboratory can grow indefinitely if plant nutrients, light, and water are provided, thus producing unlimited duckweed test specimens for use at any moment.

Duckweed is a floating organism. It is especially susceptible to surface active




FIG. 1. Quality of control tests as indicated by doubling time of duckweed frond number. Redrawn from Wang (1987a).

substances, hydrophobic (water interface (Wu *et al.*,

There are indications that it is commonly referred to reported that duckweed (*L. minor*) inhabiting a coal-ash retainir imately 95% of the surface grew under a wide range of conditions. Seto *et al.* (1979) reported Cd toxicity, however, was not observed when specimens were cultured. The plant was cultured in a high pH solution.

There are other indications of toxicity. Wang (1986a) compared duckweed and fish species sensitivity to pollutants. Of special interest was the sensitivity of duckweed and fish species to toxic substances as sensitive or sometimes more sensitive.

Duckweed is also sensitive to toxic substances. Wang (1982) reported that duckweed was more sensitive than juvenile rainbow trout, when exposed to toxic substances. Duckweed were comparable to fish in sensitivity. Wang (1988) reported that the toxicity of duckweed to *Navicula pelliculosa* was higher than those of *Navicula pelliculosa* when these species were 0.17, 0.17, and 0.17. Wang (1988) observed the toxic effects of duckweed on grass shrimp, fish, and daphnia.

The preceding discussion of duckweed plants are described in detail. On the one hand the plants are mentioned in the literature. On the other hand the plants are mentioned on the basis that the culture conditions, free of toxic substances, mental toxicity when they are cultured.

#### COMPARISON OF FISH AND DUCKWEED

Cd
Cr (VI)
Cu
Pb
Ni
Se
Zn

Note. Reproduced from Wang (1987a) Ltd.

substances, hydrophobic compounds, and the like that concentrate at the air-water interface (Wu *et al.*, 1980).

There are indications that duckweed is tolerant to environmental toxicity, and it is commonly referred to as the "carp" of plant species. Clark *et al.* (1981) reported that duckweed (*L. perpusilla*) was the most abundant macrophyte inhabiting a coal-ash retaining basin of a coal-fired power plant, occupying approximately 95% of the surface area. Gabrielson *et al.* (1980) found that duckweed grew under a wide range of nutrient conditions including high metal concentrations. Seto *et al.* (1979) reported that Cd caused chlorosis and death to *L. gibba*. Cd toxicity, however, was greatly influenced by the nutrient levels in which the specimens were cultured. Duckweed was much more tolerant to Cd when the plant was cultured in a higher-nutrient solution than in a low-nutrient solution.

There are other indications, however, suggesting that duckweed is sensitive to toxicity. Wang (1986a) conducted a series of duckweed toxicity tests on 16 aquatic pollutants. Of special interest is the comparison of the sensitivity to metal toxicity of duckweed and fish species. The results in Table 1 indicate that duckweed was as sensitive or sometimes more sensitive to metal toxicity than fish species.

Duckweed is also sensitive to organic compounds. For example, Rowe *et al.* (1982) reported that duckweed was as sensitive to epoxystearic acid toxicity as juvenile rainbow trout, while the severe toxic effects of catechol compounds on duckweed were comparable with those on daphnids. Furthermore, Hughes *et al.* (1988) reported that the toxic effects of atrazine on *L. gibba*, *Anabaena flos-aqua*, and *Navicula pelliculosa* were comparable. The 50% inhibition concentrations for these species were 0.17, 0.23, and 0.06 mg/liter, respectively. Huber *et al.* (1982) observed the toxic effects of pentachlorophenol on *L. minor* to be comparable to those on grass shrimp, fish, and snails.

The preceding discussion appears to be contradictory. On the one hand duckweed plants are described as tolerant to environmental toxicity, while on the other hand the plants are mentioned as sensitive to toxicity. The contradiction can be explained on the basis that the plants may be highly adaptive. Under optimum culture conditions, free of contamination, the plants are responsive to environmental toxicity when they encounter it. At sublethal range, the duckweed plants

TABLE 1  
COMPARISON OF FISH AND DUCKWEED BIOASSAYS ON METAL TOXICITY (mg/liter)

	Fish 96 hr LC <sub>50</sub>	Duckweed 96 hr EC <sub>50</sub>
Cd	0.92	0.2
Cr (VI)	58.5	35
Cu	0.08-1.2 (three species)	1.1
Pb	27.8	8
Ni	13.6-48.8 (four species)	0.45
Se	18.7	2.4
Zn	0.4-55 (six species)	10

Note. Reproduced from Wang (1986a) with permission from Elsevier Applied Science Publishers Ltd.

may adapt and/or develop resistance quickly due to their fast growth rate. Several reports (Duncan and Klaverkamp, 1983; Benson and Birge, 1985; Dixon and Sprague, 1981) indicate that fish can develop tolerance. Perhaps the same induced tolerance is the reason that duckweed plants are sometimes insensitive to environmental toxicity.

Duckweed is uniquely suitable for testing herbicide pollution in the aquatic environment. The use of herbicides has steadily increased in the past 25 years (Nielsen and Lee, 1987), and herbicide runoff into surface waters is widespread (Leonard, 1988). Results obtained by Taraldsen and Norberg-King (in press) demonstrated clearly that tests of duckweed, fathead minnow, and *Ceriodaphnia* complemented each other for testing different types of wastewaters. No single test is likely to be able to detect toxicity in every environmental sample.

#### Duckweed vs Algal Toxicity Tests

Both algae and higher plants are primary producers and essential parts of a balanced ecosystem. There has been a persistent question as to whether algal species can be a surrogate for plant species and whether the plant kingdom is sufficiently represented by algal species alone as a part of the three base-set tests.

In the aquatic environment, algae and duckweed are competitors. There are reports indicating that algal growth might interfere with the growth of aquatic macrophytes through alleopathic effects (Van Vierssen and Prins, 1985; Kemp *et al.*, 1984). Duckweed plants, being floating organisms, suppress algal growth by blocking sunlight.

There has been some information directly comparing results of duckweed and algal toxicity tests. As mentioned, Hughes *et al.* (1988) reported that *L. gibba*, *Anabaena*, and *Navicula* were almost equally sensitive to atrazine.

There are indications that duckweed is less sensitive to toxicity than algae. Mangi *et al.* (1978) reported that both *L. minor* and *Spirodela polyrhiza* were less sensitive to Cr (VI) than single or filamentous algae. Bioassays of solid waste leachate from coal gasifiers with *Selenastrum capricornutum* and *S. oligorhiza* showed that the alga was more sensitive to toxicity than the duckweed (Klaine, 1985). Klaine reported the 50% inhibition effect concentrations for the test species to be 55 and 76% leachate concentration, respectively.

Rowe *et al.* (1982) compared toxicity test results for spent chlorination and caustic extraction liquors using an alga *Chlorella pyrenoidesa* and a duckweed *L. perpusilla*. They found that the algae tended to be more sensitive than the duckweed to chlorinated phenolic compounds.

These results, however, should be interpreted with caution. As discussed in the following section on end points, the conventional duckweed test grossly underestimates toxic effect. The advancement of a duckweed test protocol will improve the test results in the future. Additionally, Klaine (1985) raised an important point that "Since the alga is distributed throughout the water column and the duckweed only inhabits the water surface, the alga maintains a more constant exposure to the toxicant." Similarly, results were also reported that submerged plant species generally absorbed more Cd, Cu, Pb, and Zn than the floating species, *L. gibba* and *S. polyrhiza* (Van der Werff and Pruyt, 1982). One should therefore bear in

mind the different natures of their results.

With regard to terrestrial algae complement each other as toxicants (Miller *et al.*, 1985) that "algal growth inhibitory toxicity of herbicides to terrestrial one day be extrapolated to in particular. More research and duckweed.

Unlike algal toxicity tests of effluent biomonitoring. Mar With these samples, filtration loss of sample integrity. Do sample "as is." More important wastewaters are labile because These samples require either tested in either manner, which

Duckweed toxicity tests compared algal tests. Nasu *et al.* (1984) cation and frond growth (while the Cd ion suppressed growth. Interesting findings from studies of environmental toxic

DUC

#### Species

Many duckweed species have been used in toxicity tests. *L. polyrhiza*, *L. perpusilla*, and *S. oligorhiza*. Features of three species

There are relatively few data available. Coley (1985) compared the toxicity of duckweed to aqueous extracts of natural materials. He reported that *L. gibba* was the most sensitive to the extract of coal distillate compared to *L. perpusilla*, whereas complete inhibition was observed for *S. oligorhiza*.

Under low irradiance, Toxicity was most pronounced for *L. gibba* and *S. oligorhiza* compared to *L. valdiviana*. *S. oligorhiza* was least tolerant to Cr, while *L. gibba* was most tolerant to concentrations above 10 mg/liter.

The selection of test species for toxicity to the toxicant, historical data, and comparability, two

mind the different natures of the tests for algae and duckweed when comparing their results.

With regard to terrestrial plant species, it is apparent that higher plants and algae complement each other as toxicity test species when exposed to varieties of toxicants (Miller *et al.*, 1985; Thomas *et al.*, 1986). Garten and Frank (1984) stated that "algal growth inhibition tests cannot be used generally to predict phytotoxicity of herbicides to terrestrial plant species." Perhaps the same conclusion may one day be extrapolated to include aquatic macrophytes in general and duckweed in particular. More research is needed for comparative toxicity tests using algae and duckweed.

Unlike algal toxicity tests, duckweed toxicity tests are especially suitable for effluent biomonitoring. Many industrial and municipal wastewaters are turbid. With these samples, filtration is required to conduct algal tests, resulting in the loss of sample integrity. Duckweed tests, however, can be performed on the sample "as is." More importantly, some industrial and almost all municipal wastewaters are labile because of high organic content and microbial population. These samples require either flow-through or renewal tests. Duckweed can be tested in either manner, while algae cannot.

Duckweed toxicity tests can also reveal effects that cannot be obtained by using algal tests. Nasu *et al.* (1984) observed that Cu suppressed both frond multiplication and frond growth (wet weight increase of each frond) of *L. paucicostata*, while the Cd ion suppressed only frond multiplication and did not affect frond growth. Interesting findings such as these are very important for comparative studies of environmental toxicology.

good duckweed plants

## DUCKWEED TOXICITY TESTS


### Species

Many duckweed species have been studied: *L. minor*, *L. gibba*, *L. valdiviana*, *L. polyrrhiza*, *L. perpusilla*, *L. paucicostata*, *S. polyrhiza*, *S. punctata*, and *S. oligorhiza*. Features of three common duckweed species are depicted in Fig. 2.

There are relatively few data for comparative studies on different species. King and Coley (1985) compared the sensitivity of *L. gibba*, *L. minor*, and *L. perpusilla* to aqueous extracts of natural and synthetic oils, as well as to coal distillate. They reported that *L. gibba* was the most resistant among the three species. The 10% extract of coal distillate completely inhibited the growth of *L. minor* and *L. perpusilla*, whereas complete inhibition of *L. gibba* was noted only in the 15% extract.

Under low irradiance, Takemoto and Noble (1986) found that sulfite toxicity was most pronounced for *L. gibba*, less marked for *S. oligorhiza*, and not observed with *L. valdiviana*. Staves and Knaus (1985) reported that *S. polyrhiza* was least tolerant to Cr, while *S. punctata* and *L. gibba* were more tolerant to concentrations above 10 mg/liter.

The selection of test species is usually based on specimen availability, sensitivity to the toxicant, historical data, and the like. To encourage test standardization and comparability, two species have been recommended. The American



*Spirodela polyrhiza*, x 2.4      *Lemna minor*, x 2.4      *Lemna perpusilla*, x 2.4

FIG. 2. Features of *Spirodela polyrhiza*, *Lemna minor*, and *Lemna perpusilla*. Reproduced from Correll and Correll (1972).

Society for Testing and Materials (in press) recommends *L. gibba*, while *Standard Methods for Examination of Water and Wastewater* in its 17th Edition Supplement (American Public Health Association *et al.*, in press) endorses *L. minor*. Although the use of standard species is encouraged, other species should be tested so that new information may help to evolve test methods in the future.

#### Test Types

Duckweed toxicity tests can be used in static, renewal, or flow-through experiments. Static experiments are simple and economical. They are especially useful for screening tests of unknown samples or samples which contain toxic metals (Wang, 1986a, b, 1987a, b, 1988; Wang and Williams, 1988). Typical test conditions for conducting duckweed static experiment are given in Table 2. Flow-through and renewal experiments have also been reported (Bishop and Perry,

TABLE 2  
TYPICAL DUCKWEED STATIC TEST

Temperature	27–28°C
Light quality	Cool white fluorescent
Light intensity	86 $\mu\text{E}/\text{m}^2/\text{sec}$
Photoperiod	Continuous
Test vessel	60 × 15-mm culture dish
Test solution/vessel	15 ml
Test specimens/vessel	20 fronds (10 colonies)
Replicates	4
Water control and dilution water	Duckweed nutrient medium, i.e., 10-fold algal nutrient medium
Test duration	120 hr
End point	Frond increase/vessel

Note. Modified from *Standard Methods* (American Public Health Association *et al.*, in press).

1981; Davis, 1981; Walbiments are useful for sa

#### Test Vessel

Several different test tubes, jars, Erlenmeyer Kugimoto, 1981; Hutch Wang, 1986a, b). In gen experiment. Plastic vess duckweed plants to the excessive evaporation c

The test solution in a more to accommodate d can be used. Recently tl 15-mm culture dishes. were placed in each disl nutrient solution (Amer 150-ml solution contain iaturization of the test frond doubling time of c with a total of 22 tests.  $1.9 \pm 0.36$  days as show a 30-ml plastic cup con

The advantage of min more flexible in the expe test volume also minin pleted.

#### Control Tests

There are two types o Almost all toxicity tests The negative control se reference point to which from time to time as sh

Positive controls are as negative controls. A is added to determine t compounds have been tachlorophenol, phenol chromate (Davis and F Threader and Houston, growth of *L. minor* unc duckweed response to toxicity to *L. minor* wa

1981; Davis, 1981; Walbridge, 1977). In general, flow-through and renewal experiments are useful for samples containing volatile or biodegradable compounds.

### *Test Vessel*

Several different test vessels have been used: glass beakers, flat-bottomed test tubes, jars, Erlenmeyer flasks, and culture dishes (Fekete *et al.*, 1976; Nasu and Kugimoto, 1981; Hutchinson and Czyska, 1975; Stanley and Madewell, 1976; Wang, 1986a, b). In general, only one type of test vessel should be used in an experiment. Plastic vessels are not recommended because of strong adherence of duckweed plants to the plastic walls. The test vessel should be covered to avoid excessive evaporation of water and test sample.

The test solution in a vessel in general should have a liquid height of 5 cm or more to accommodate duckweed root length. Either 100 or 200 ml of test solution can be used. Recently the author reduced the test solution to 15 ml by using 60 × 15-mm culture dishes. The liquid depth was 5 mm. Twenty fronds of *L. minor* were placed in each dish. *Lemna* grew in a 15-ml solution containing 10-fold algal nutrient solution (American Public Health Association *et al.*, 1985) as well as in a 150-ml solution containing 2-fold algal nutrient solution (Wang, 1987a). The miniaturization of the test solution gave reproducible results, as evidenced by the frond doubling time of control samples  $1.6 \pm 0.13$  days in a 6-month test period, with a total of 22 tests. This result compared favorably with the doubling time of  $1.9 \pm 0.36$  days as shown in Fig. 1. Taraldsen and Norberg-King (in press) used a 30-ml plastic cup containing 15 ml of test solution.

The advantage of miniaturizing the test solution is obvious. In addition to being more flexible in the experimental design because of its smaller volume, the smaller test volume also minimizes disposal problems after the experiments are completed.

### *Control Tests*

There are two types of control samples: negative controls and positive controls. Almost all toxicity tests use a negative control where no test substance is present. The negative control serves as a quality control in an experiment as well as a reference point to which a test sample is compared. Negative control values vary from time to time as shown in Fig. 1.

Positive controls are less frequently employed, although they are as important as negative controls. A positive control is a sample to which a reference toxicant is added to determine the degree of response over time. Organic and inorganic compounds have been studied as potential reference toxicants, including pentachlorophenol, phenol, sodium dodecyl sulfate, sodium chloride, cadmium, and chromate (Davis and Hoos, 1975; Fogels and Sprague, 1977; Jop *et al.*, 1986; Threader and Houston, 1983). Wang (1987a) found that in an 18-month period, the growth of *L. minor* underwent cyclic changes in the negative control, while the duckweed response to Cr toxicity was nearly constant (Fig. 3). In general, Cr toxicity to *L. minor* was not ameliorated by differences in water quality factors




FIG. 3. Duckweed growth in water control (■) and in sample control (□), and Cr toxicity in nutrient medium (●) and in enriched water samples (○), using Illinois River samples. Redrawn from Wang (1987a).

such as suspended solids or dissolved fraction (Wang, 1986b). On the basis of these and other results, Cr (VI) has been recommended as a universal reference toxicant.

#### Nutrient Media

Duckweed toxicity tests, which generally rely on growth and multiplication as the test end point, require sufficient plant nutrients for optimum conditions. Plant nutrients are uniformly added to control and test samples.

Several nutrient media have been reported. They included the medium used by Ballard (1966), Bristol's medium as given by Eyster (1968); Jacob's medium as given by McLay (1976); Hoagland's, Hunter's, and Bonner-Devirian's media as given by Nasu and Kugimoto (1981); and the medium used by Fekete *et al.* (1976). The composition varied widely among these media.

Nasu and Kugimoto (1981) found that the pH of the nutrient media, the concentration and composition of the nutrients in the medium, and the temperature at which cultures were maintained affected the sensitivity of *L. paucicostata* to heavy metals. They recommended the use of Bonner-Devirian's medium at temperatures above 25°C. Nasu *et al.* (1983) reported that the absorption of Cu and Cd in *L. paucicostata* was suppressed by the addition of EDTA to the medium. Only 30  $\mu\text{M}$  of EDTA was sufficient to prevent Cu absorption at the concentration of 5–10  $\mu\text{M}$ , whereas 400  $\mu\text{M}$  of EDTA was required to prevent Cd absorption at the same concentration as Cu. They found that the growth inhibition of *Lemna* was proportional to the amount of metal absorption.

The green alga *Selenastrum capricornutum* and the common duckweed *L. mi-*

nor are frequently used for share the same nutrient stock double-strength algal nutrient solutions recommended by The same approach has been Materials (in press), who will also be recommended in *et al.*, in press) who suggested.

#### End Points

Many end points have been points are generally based on plant number, root number, C-14 uptake, chlorophyll, and Culley *et al.*, 1981; Lockha McNabb, 1978; Sahai *et al.*

The most commonly used end point is bud inclusion in order to be repeatedly until accurate results are obtained. These end points are nondestructive. These end points are:

The first disadvantage is that it has been observed frequently.

PREPARATION

Stock
A
B
C

Note. Ten milliliters each of three (10 $\times$ ) of algal nutrient solution. (A)

nor are frequently used for phytotoxicity testing. It would be highly desirable to share the same nutrient stock solutions for these two tests. Wang (1986c) used double-strength algal nutrient solution, which is recommended by *Standard Methods* (American Public Health Association *et al.*, 1985). Instead of seven stock solutions recommended by *Standard Methods*, Wang prepared three solutions. The same approach has been adopted by the American Society for Testing and Materials (in press), who recommend 20-fold (20×) strength. This same approach will also be recommended in *Standard Methods* (American Public Health Association *et al.*, in press) where 10-fold (10×) strength as shown in Table 3 will be suggested.

#### End Points

Many end points have been used to express duckweed test results. These end points are generally based on the population of duckweed plants: frond number, plant number, root number, dry or fresh biomass, root length, frond diameter, C-14 uptake, chlorophyll, and the like (Bishop and Perry, 1981; Said *et al.*, 1979; Culley *et al.*, 1981; Lockhart *et al.*, 1983; Bishop and Perry, 1981; Glandon and McNabb, 1978; Sahai *et al.*, 1977; Fekete *et al.*, 1976).

The most commonly used test end point is frond number. Any visible, protruding bud is included in order to avoid individual bias. The frond count can be made repeatedly until accurate results are obtained. This determination is simple, rapid, and nondestructive. These advantages, however, are weighted against disadvantages.

The first disadvantage is that frond count is irrelevant to frond size or biomass. It has been observed frequently that under toxic stress small buds may protrude

TABLE 3  
PREPARATION OF DUCKWEED NUTRIENT SOLUTION

Stock	Compound	Concentration (mg/liter)
A	NaNO <sub>3</sub>	22,500
	NaHCO <sub>3</sub>	15,000
	K <sub>2</sub> HPO <sub>4</sub>	1,040
B	MgCl <sub>2</sub> · 6H <sub>2</sub> O	12,164
	CaCl <sub>2</sub> · 2H <sub>2</sub> O	4,410
	FeCl <sub>3</sub>	96
	Na <sub>2</sub> EDTA · 2H <sub>2</sub> O	300
	MnCl <sub>2</sub>	264
C	MgSO <sub>4</sub> · 7H <sub>2</sub> O	14,700
	H <sub>3</sub> BO <sub>3</sub>	185.5
	ZnCl <sub>2</sub>	3.270
	CoCl <sub>2</sub>	0.780
	NaMoO <sub>4</sub> · 2H <sub>2</sub> O	7.260
	CuCl <sub>2</sub>	0.009

Note. Ten milliliters each of three stock solutions are diluted into 1 liter to make 10-fold strength (10×) of algal nutrient solution. (American Public Health Association *et al.*, in press).

and be counted as individual fronds. A small bud may be less than 5% of the biomass of a healthy frond in a control group, yet these two fronds are considered equal. The result of using frond count as the end point is to grossly underestimate the toxic effect to duckweed. In comparison, algal response to toxicity is unlikely to exhibit such a distortion. This is a reason why results of duckweed and algal toxicity tests should be interpreted with caution.

The other and more important disadvantage is that frond count does not reflect whether the plant is alive or dead. This problem was clearly demonstrated in an experiment using industrial effluents (Wang and Williams, 1988). In the effluent, new fronds developed and died during 96 hr exposure. Live and dead fronds could not be distinguished by using the frond count method.

Cowgill and Milazzo (1989) indicated that determinations of dry biomass (constant weight at 60°C) were the least time-consuming and least subject to human error. This method, although an improvement by taking into account change in biomass as a toxic response, does not distinguish between live and dead plants.

Duckweed plants can also exhibit many symptoms when they are under stress. These symptoms include chlorosis (loss of pigment), necrosis (localized dead tissue), colony breakup, root destruction, loss of buoyancy, and gibbosity (humpback or swelling). For example, Cd is known to cause chlorosis (Seto *et al.*, 1979).

Potentially the most sensitive and accurate test end points include viable biomass and physiological activity. Examples are tests of adenosine triphosphate content, fluorescence emission (Caux *et al.*, 1988), C-14 uptake (Lockhart *et al.*, 1983), and oxygen production (Huber *et al.*, 1982). More studies are needed in this area.

### Test Results

Numerous duckweed toxicity tests have been conducted. Table 4 gives test results for single compounds, including organic and inorganic substances. Table 5 shows the test results for complex mixtures.

### Applications

Duckweed toxicity tests are highly versatile in the aquatic environment. The tests are applicable to lake, river, or ground water; single chemical compounds or complex effluents from industrial or municipal sources; organic or inorganic compounds; rain samples; and sediment samples (Wang, 1986a-d, 1987a, b; Wang and Williams, 1988, 1990; Hartman and Martin, 1984; Fekete *et al.*, 1976). Wang and Williams (1988, 1990) reported that duckweed was more sensitive to industrial effluents than higher plants such as cabbage and millet.

Sediment toxicity testing using duckweed is especially noteworthy because sediment samples can be tested either as aqueous extract or as sediment slurry (Wang, 1986d). Some compounds remained biologically active in the presence of soil particles. Birmingham and Colman (1983) found that both *Lemna* species and an alga, *Anabaena flos-aquae*, detected significant phytotoxicity of a herbicide, Reglone A-R, in a soil-water mixture at a 33-ppm concentration level. Hartman and Martin (1985) also reported that the presence of suspended sediment had a minor effect on the toxicity of alachlor and atrazine to *L. minor*.

Test substances	Spe
Ba	<i>Lemna n</i>
Ba	<i>L. minor</i>
B	<i>L. minor</i>
Cd	<i>L. minor</i>
Cd	<i>L. valdiv</i>
Cd	<i>L. valdiv</i>
Cd	<i>L. valdiv</i>
CdCl <sub>2</sub>	<i>L. polyrr</i>
CdSO <sub>4</sub>	<i>L. polyrr</i>
CdCl <sub>2</sub>	<i>L. pauci</i>
Cl	<i>L. minor</i>
Cr(VI)	<i>L. minor</i>
Cr(VI)	<i>L. minor</i>
Cr(VI)	<i>Spirodei</i>
Cu	<i>L. mino</i>
Cu	<i>L. valdi</i>
Cu	<i>L. mino</i>
Cu	<i>L. mino</i>
CuSO <sub>4</sub>	<i>L. gibb</i>
CuSO <sub>4</sub>	<i>L. gibb</i>
Pb	<i>L. mino</i>
Mn	<i>L. minc</i>
Ni	<i>L. minc</i>
Ni	<i>L. minc</i>
Se	<i>L. minc</i>
SO <sub>4</sub>	<i>L. minc</i>
Zn	<i>L. minc</i>
Zn	<i>L. minc</i>
AE	<i>L. min</i>
Alachlor	<i>L. min</i>
Atrazine	<i>L. min</i>
Carbofuran	<i>L. min</i>
O-Cresol	<i>L. gibb</i>
O-Cresol	<i>L. gibb</i>
CTAC	<i>L. min</i>
DEP	<i>L. gibb</i>
EG	<i>L. gibb</i>
Glyphosate	<i>L. min</i>
LAS	<i>L. min</i>
Pentachlorophenol	<i>L. min</i>
Phenol	<i>L. min</i>
Salicylic acid	<i>L. min</i>
SDS	<i>L. min</i>
2,4,6-TCP	<i>L. gibb</i>
2,4,6-TCP	<i>L. gibb</i>

Note. All IC<sub>50</sub>'s (50% inhibit increase, except as noted. AE (2-ethylhexyl)phthlate; EG, Eth 2,4,6-TCP, 2,4,6-trichlorophenol  
<sup>a</sup> Flow-through test.  
<sup>b</sup> Oxygen production as test end point.  
<sup>c</sup> Growth rate as test end point.

The use of herbicide tests in the aquatic environment. Herbicide tests are an important part of the duckweed tests.

TABLE 4  
DUCKWEED TOXICITY TESTS OF SINGLE COMPOUNDS

Test substances	Species	Duration	IC <sub>50</sub> 's	References
Ba	<i>Lemna minor</i>	4 days	24 mg/liter	Wang (1988)
Ba	<i>L. minor</i>	4	26	Wang (1986a)
B	<i>L. minor</i>	4	>60	Wang (1986a)
Cd	<i>L. minor</i>	4	0.2	Wang (1986a)
Cd	<i>L. valdiviana</i>	7	0.15	Hutchinson and Czyrska (1975)
Cd	<i>L. valdiviana</i>	14	0.3	Hutchinson and Czyrska (1975)
Cd	<i>L. valdiviana</i>	21	0.3	Hutchinson and Czyrska (1975)
CdCl <sub>2</sub>	<i>L. polyrrhiza</i>	14	0.9	Charpentier <i>et al.</i> (1987)
CdSO <sub>4</sub>	<i>L. polyrrhiza</i>	14	0.6	Charpentier <i>et al.</i> (1987)
CdCl <sub>2</sub>	<i>L. paucicostata</i>	7	10	Nasu <i>et al.</i> (1984)
Cl	<i>L. minor</i>	4	930	Wang (1986a)
Cr(VI)	<i>L. minor</i>	4	35	Wang (1986a)
Cr(VI)	<i>L. minor</i>	14	6	Mangi <i>et al.</i> (1978)
Cr(VI)	<i>Spirodela polyrrhiza</i>	14	>10	Mangi <i>et al.</i> (1978)
Cu	<i>L. minor</i>	4	1.1	Wang (1986a)
Cu	<i>L. valdiviana</i>	21	0.14	Hutchinson and Czyrska (1975)
Cu	<i>L. minor</i>	7 <sup>a</sup>	0.119	Walbridge (1977)
Cu	<i>L. minor</i>	7 <sup>a</sup>	0.8	Bishop and Perry (1981)
CuSO <sub>4</sub>	<i>L. gibba</i>	7 <sup>c</sup>	2.21	Davis (1981)
CuSO <sub>4</sub>	<i>L. gibba</i>	7 <sup>a,c</sup>	3.51	Davis (1981)
Pb	<i>L. minor</i>	4	8	Wang (1986a)
Mn	<i>L. minor</i>	4	31	Wang (1986a)
Ni	<i>L. minor</i>	4	0.45	Wang (1986a)
Ni	<i>L. minor</i>	4	0.30	Wang (1987b)
Se	<i>L. minor</i>	4	2.4	Wang (1986a)
SO <sub>4</sub>	<i>L. minor</i>	4	>1,000	Wang (1986a)
Zn	<i>L. minor</i>	4	10	Wang (1986a)
AE	<i>L. minor</i>	7 <sup>a</sup>	21	Bishop and Perry (1981)
Alachlor	<i>L. minor</i>	14	0.01	Hartman and Martin (1985)
Atrazine	<i>L. minor</i>	14	>0.1	Hartman and Martin (1985)
Carbofuran	<i>L. minor</i>	14	>10	Hartman and Martin (1985)
O-Cresol	<i>L. gibba</i>	7 <sup>a,c</sup>	152	Davis (1981)
O-Cresol	<i>L. gibba</i>	7 <sup>c</sup>	246	Davis (1981)
CTAC	<i>L. minor</i>	7 <sup>a</sup>	0.1	Bishop and Perry (1981)
DEP	<i>L. gibba</i>	7 <sup>c</sup>	2,060	Davis (1981)
EG	<i>L. gibba</i>	7 <sup>c</sup>	17,159	Davis (1981)
Glyphosate	<i>L. minor</i>	7	2	Hartman and Martin (1984)
LAS	<i>L. minor</i>	7 <sup>a</sup>	2.7	Bishop and Perry (1981)
Pentachlorophenol	<i>L. minor</i>	60 hr	3.2 <sup>b</sup>	Huber <i>et al.</i> (1982)
Phenol	<i>L. minor</i>	4	>12	Wang (1986a)
Salicylic acid	<i>L. minor</i>	7	107	Wang and Lay (1989)
SDS	<i>L. minor</i>	7 <sup>a</sup>	43	Bishop and Perry (1981)
2,4,6-TCP	<i>L. gibba</i>	7 <sup>a,c</sup>	0.03	Davis (1981)
2,4,6-TCP	<i>L. gibba</i>	7 <sup>c</sup>	0.13	Davis (1981)

Note. All IC<sub>50</sub>'s (50% inhibition concentrations) are obtained with static tests and the test end point is frond count increase, except as noted. AE, C<sub>14.5</sub> alcohol ethoxylate; CATC, Cetyl trimethyl ammonium chloride; DEP, Di-(2-ethylhexyl)phthalate; EG, Ethylene glycol; LAS, C<sub>11.8</sub> linear alkylbenzene sulfonate; SDS, Sodium dodecyl sulfate; 2,4,6-TCP, 2,4,6-trichlorophenol.

<sup>a</sup> Flow-through test.

<sup>b</sup> Oxygen production as test end point.

<sup>c</sup> Growth rate as test end point.

The use of herbicides in the United States increased 280% between 1966 and 1981 (Nielsen and Lee, 1987) and resulted in their widespread distribution in the environment. Herbicides such as alachlor, atrazine, butylate, and cyanazine have been detected in fog and rain (Richards *et al.*, 1987; Glotfelty *et al.*, 1987). Duckweed tests are an inexpensive and sensitive biomonitoring tool for detecting phy-

TABLE 5  
DUCKWEED TOXICITY TESTS OF COMPLEX SAMPLES

Sample	Species	Duration	IC <sub>50</sub>	References
SRC-11 fuel oil	<i>L. perpusilla</i>	8 days	3.4%	King and Coley (1985)
SRC-11 fuel oil	<i>L. minor</i>	8	4	King and Coley (1985)
SRC-11 fuel oil	<i>L. gibba</i>	8	4.2	King and Coley (1985)
Raw coal distillate	<i>L. minor</i>	8	3.5	King and Coley (1985)
Raw coal distillate	<i>L. perpusilla</i>	8	3.8	King and Coley (1985)
Raw coal distillate	<i>L. gibba</i>	8	4.2	King and Coley (1985)
Coal ash leachate	<i>S. oligorhiza</i>	NA	76	Klaine (1985)
PIE 1	<i>L. minor</i>	5	20	Wang and Williams (1988)
2	<i>L. minor</i>	5	<2	Wang and Williams (1988)
PIE 1	<i>L. minor</i>	4	38	Wang and Williams (1990)
2	<i>L. minor</i>	4	49	Wang and Williams (1990)
3	<i>L. minor</i>	4	49	Wang and Williams (1990)
4	<i>L. minor</i>	4	22	Wang and Williams (1990)
5	<i>L. minor</i>	4	29	Wang and Williams (1990)
6	<i>L. minor</i>	4	91	Wang and Williams (1990)
7	<i>L. minor</i>	4	43	Wang and Williams (1990)
8	<i>L. minor</i>	4	<1.6	Wang and Williams (1990)
9	<i>L. minor</i>	4	<1.6	Wang and Williams (1990)
PIE 1	<i>L. minor</i>	5	73	Wang (unpublished)
2	<i>L. minor</i>	5	47	Wang (unpublished)

Note. All IC<sub>50</sub>'s (50% inhibition concentration, %) are obtained with static tests and the test end point is frond count increase. NA, not available; PIE, pretreated industrial effluent.

tototoxicity in fog and rain samples. Effects of acid rain can also be measured by using duckweed (Schindler, 1988).

#### REFERENCES

- American Public Health Association, American Water Works Association, and Water Pollution Control Federation (1985). "Standard Methods for Examination of Water and Wastewater." Washington, DC.
- American Public Health Association, American Water Works Association, and Water Pollution Control Federation, in press. Section 8211 toxicity test procedures using common duckweed, *Lemna minor* (proposed). In "Standard Methods for Examination of Water and Wastewater," 17th ed. Suppl., Washington, DC.
- American Society for Testing and Materials, in press. Proposed new standard guide for conducting static toxicity tests with *Lemna gibba* G3.
- Ballard, E. G. (1966). A comparison study of organic nitrogenous compounds to serve as sole sources of nitrogen for growth of plants. *Plant Soil* 25, 153-166.
- Benson, W. H., and Birge, W. J. (1985). Heavy metal tolerance and metallothionein induction in fathead minnows: Results from field and laboratory experiments. *Environ. Toxicol. Chem.* 4, 209-217.
- Birmingham, B. C., and Colman, B. (1983). Potential phytotoxicity of diquat accumulated by aquatic plants and sediments. *Water, Air, Soil Pollut.* 19, 123-131.
- Bishop, W. E., and Perry, R. L. (1981). Development and evaluation of a flowthrough growth inhibition test with duckweed (*Lemna minor*). ASTM STP 737, pp. 421-435. Amer. Soc. Testing Materials.
- Caux, P. Y., Weinberger, P., and Carlisle, D. B. (1988). A physiological study of the effects of Triton surfactants on *Lemna minor* L. *Environ. Toxicol. Chem.* 7, 671-676.
- Charpentier, S., Garnier, J., and Fl. experimental cultures of duckweed 1055-1061.
- Clark, J. R., Van Hassel, V. H., Nic and depuration of metals by duckweed 1055-1061.
- Correll, D. S., and Correll, H. B. (1985). EPA Water Pollution Control 1 Washington, DC.
- Cowgill, U. M., and Milazzo, D. F. ASTM STP 1027, pp. 379-391.
- Culley, D. D., Rejmankova, E., Ke of duckweeds in aquaculture, 27-49.
- Davis, J. A. (1981). Comparison of *Lemna gibba* G3, EPA 560/6-8
- Davis, J. C., and Hoos, R. A. W. (1985). as reference toxicants for salmonids. *Environ. Toxicol. Chem.* 4, 128-138.
- Dixon, D. G., and Sprague, J. B. (1985). modifying factor in toxicity. *Environ. Toxicol. Chem.* 4, 128-138.
- Duncan, D. A., and Klaverekamp, (1985). (*Calostomus commersoni*) prey. *Fish. Aq. Sci.* 40, 128-138.
- Eyster, C. (1968). Microorganic ar Environment," (D. Jackson, F. Fekete, A., Riemer, D. N., and M. the release of available phosphorus. *Environ. Toxicol. Chem.* 4, 128-138.
- Fogels, A., and Sprague, J. B. (1985). rainbow trout to five poisons. *Environ. Toxicol. Chem.* 4, 128-138.
- Frick, H. (1985). Micronutrient tol 8, 1131-1145.
- Gabrielson, F. C., Malatino, A. M. in phosphorus and nitrogen sp 602-605.
- Garten, C. T., and Frank, M. L. (1985). inhibition by herbicides. ORN Oak Ridge National Laborato 53-64.
- Glandon, R. P., and McNabb, C. 53-64.
- Glotfelty, D. E., Seiber, J. N., ar 602-605.
- Hartman, W. A., and Martin, D. of glyphosate to *Daphnia pul.*
- Hartman, W. A., and Martin, D. *Lemna minor*, and *Potamogeton*
- Hillman, W. S. (1961). The Lemn literature. *Botan. Rev.* 27, 22
- Hillman, W. S., and Culley, D. I. toxicity of effluents and receiving water. Environmental Protection Agency, C Huber, W., Schubert, V., and S. the aquatic macrophyte *Lemna*
- Hughes, J. S., Alexander, M. M. toxicity in aquatic plant bioassays. *Environ. Toxicol. Chem.* 7, 671-676.

- entier, S., Garnier, J., and Flaugnatti, R. (1987). Toxicity and bioaccumulation of cadmium in experimental cultures of duckweed, *Lemna polyrrhiza* L. *Bull. Environ. Contam. Toxicol.* 38, 1055-1061.
- Clark, J. R., Van Hassel, V. H., Nicholson, R. B., Cherry, D. S., and Cairns, J. (1981). Accumulation and depuration of metals by duckweed (*Lemna perpusilla*). *Ecotoxicol. Environ. Saf.* 5, 87-96.
- Correll, D. S., and Correll, H. B. (1972). Aquatic and wetland plants for Southwestern United States. EPA Water Pollution Control Research Series 16030 DNL. Environmental Protection Agency, Washington, DC.
- Cowgill, U. M., and Milazzo, D. P. (1989). The culturing and testing of two species of duckweed. ASTM STP 1027, pp. 379-391. Amer. Soc. Testing Materials.
- Culley, D. D., Rejmankova, E., Kevt, J., and Frye, J. B. (1981). Production, chemical quality and use of duckweeds in aquaculture, waste management and animal feeds. *J. World Maric. Soc.* 12, 27-49.
- Davis, J. A. (1981). Comparison of static-replacement and flow-through bioassays using duckweed, *Lemna gibba* G3, EPA 560/6-81-003. U.S. Environmental Protection Agency, Washington, D.C.
- Davis, J. C., and Hoos, R. A. W. (1975). Use of sodium pentachlorophenate and dehydroabietic acid as reference toxicants for salmonid bioassays. *J. Fish. Res. Board Canad.* 32, 411-416.
- Dixon, D. G., and Sprague, J. B. (1981). Acclimation to copper by rainbow trout (*Salmo gairdneri*)—a modifying factor in toxicity. *Canad. J. Fish. Aquat. Sci.* 38, 880-888.
- Duncan, D. A., and Klaverkamp, J. F. (1983). Tolerance and resistance to cadmium in white sucker (*Catostomus commersoni*) previously exposed to cadmium, mercury, zinc, or selenium. *Canad. J. Fish. Aq. Sci.* 40, 128-138.
- Eyster, C. (1968). Microorganic and microinorganic requirements for algae. In "Algae, Man and the Environment," (D. Jackson, Ed.), pp. 27-36. Syracuse Univ. Press, Syracuse, NY.
- Fekete, A., Riemer, D. N., and Motto, H. L. (1976). A bioassay using common duckweed to evaluate the release of available phosphorus from pond sediments. *J. Aq. Plant Manag.* 14, 19-25.
- Fogels, A., and Sprague, J. B. (1977). Comparative short-term tolerance of zebrafish, flagfish, and rainbow trout to five poisons including potential reference toxicants. *Water Res.* 11, 811-817.
- Frick, H. (1985). Micronutrient tolerance and accumulation in the duckweed, *Lemna*. *J. Plant Nutrit.* 8, 1131-1145.
- Gabrielson, F. C., Malatino, A. M., and Santa Cruz, G. J. (1980). Correlation of seasonal variations in phosphorus and nitrogen species in Upper Warrior River with duckweed. NTIS NP-2903928.
- Garten, C. T., and Frank, M. L. (1984). Comparison of toxicity to terrestrial plants with algal growth inhibition by herbicides. ORNL/TM-9177, Environmental Sciences Division Publication No. 2361. Oak Ridge National Laboratory, Oak Ridge, TN.
- Glandon, R. P., and McNabb, C. D. (1978). The uptake of boron by *Lemna minor*. *Aquat. Bot.* 4, 53-64.
- Glotfelty, D. E., Seiber, J. N., and Liljedahl, L. A. (1987). Pesticides in fog. *Nature (London)* 325, 602-605.
- Hartman, W. A., and Martin, D. B. (1984). Effect of suspended bentonite clay on the acute toxicity of glyphosate to *Daphnia pulex* and *Lemna minor*. *Bull. Environ. Contam. Toxicol.* 33, 355-361.
- Hartman, W. A., and Martin, D. B. (1985). Effects of four agricultural pesticides on *Daphnia pulex*, *Lemna minor*, and *Potamogeton pectinatus*. *Bull. Environ. Contam. Toxicol.* 35, 646-651.
- Hillman, W. S. (1961). The Lemnaceae, or duckweed. A review of the descriptive and experimental literature. *Botan. Rev.* 27, 221-287.
- Hillman, W. S., and Culley, D. D. (1978). The use of duckweed. *Amer. Sci.* 66, 442-451.
- Horning, W. B., and Weber, C. I. (Eds.) (1985). Short-term methods for estimating the chronic toxicity of effluents and receiving waters to freshwater organisms. EPA/600/4-85/014. U.S. Environmental Protection Agency, Cincinnati, OH.
- Huber, W., Schubert, V., and Sauter, C. (1982). Effects of pentachlorophenol on the metabolism of the aquatic macrophyte *Lemna minor* L. *Environ. Pollut. (Ser. A)* 29, 215-223.
- Hughes, J. S., Alexander, M. M., and Balu, K. (1988). An evaluation of appropriate expressions of toxicity in aquatic plant bioassays as demonstrated by the effects of atrazine on algae and duckweed. ASTM STP 921, pp. 531-547. Amer. Soc. Testing & Materials.

- Hutchinson, T. C., and Czyska, H. (1975). Heavy metal toxicity and synergism to floating aquatic weeds. *Verh.-Int. Ver. Limnol.* 19, 2102-2111.
- Jop, K. M., Rodgers, J. H., Dorn, P. B., and Dickson, K. L. (1986). Use of hexavalent chromium as a reference toxicant in aquatic toxicity tests. ASTM STP 921, pp. 390-403. Amer. Society Testing Materials.
- Kemp, W. M., Boynton, W. R., Stevenson, J. C., Means, J. C., Twilley, R. R., and Jones, T. W. (1984). Submerged vegetation in Upper Chesapeake Bay: Studies related to possible causes of the recent decline in abundance. EPA/600/S3-84-15, NTIS PB-84-140 292.
- King, J. M., and Coley, K. S. (1985). Toxicity of aqueous extracts of natural and synthetic oils to three species of *Lemna*. ASTM STP891, pp. 302-309. Amer. Soc. Testing Materials.
- † Klaine, S. J. (1985). Toxicity of coal gasifier solid waste to the aquatic plants *Selenastrum capricornutum* and *Spirodela oligorhiza*. *Bull. Environ. Contam. Toxicol.* 35, 551-555.
- Leonard, R. A. (1988). Herbicides in surface waters. In "Environmental Chemistry of Herbicides" (R. Grover, Ed.), CRC Press, Boca Raton, FL.
- Lewis, W. M., and Bender, M. (1961). Effect of a cover of duckweed and the alga *Pithophora* upon the dissolved oxygen and free carbon dioxide of small ponds. *Ecology* 42, 602-603.
- Lockhart, W. L., Billeck, B. N., deMarch, B. E. G., and Muir, D. C. G. (1983). Uptake and toxicity of organic compounds: Studies with an aquatic macrophyte, duckweed (*Lemna minor*). ASTM STP 802, pp. 460-468. Amer. Soc. Testing Materials.
- Mangi, J., Schmidt, K., Pankow, J., Gaines, L., and Turner, P. (1978). Effects of chromium on some aquatic plants. *Environ. Pollut.* 16, 285-291.
- McLay, C. L. (1976). The effect of pH on the population growth of three species of duckweed: *Spirodela oligorhiza*, *Lemna minor*, and *Wolffia arrhiza*. *Freshwater Biol.* 6, 125-136.
- Miller, W. E., Peterson, S. A., Greene, J. C., and Callahan, C. A. (1985). Comparative toxicology of laboratory organisms for assessing hazardous waste sites. *J. Environ. Qual.* 14, 569-574.
- Nasu, Y., and Kugimoto, M. (1981). *Lemna* (duckweed) as an indicator of water pollution. I. The sensitivity of *Lemna paucicostata* to heavy metals. *Arch. Environ. Contam. Toxicol.* 10, 159-169.
- Nasu, Y., Kugimoto, M., Tanaka, O., and Takimoto, A. (1983). Comparative studies on the absorption of cadmium and copper in *Lemna paucicostata*. *Environ. Pollut. (Ser. A)* 32, 201-209.
- Nasu, Y., Kugimoto, M., Tanaka, O., Yanase, D., and Takimoto, A. (1984). Effects of cadmium and copper co-existing in the medium on the growth and flowering of *Lemna paucicostata* in relation to their absorption. *Environ. Pollut. (Ser. A)* 33, 267-274.
- Nielsen, E. G., and Lee, L. K. (1987). The magnitude and costs of groundwater contamination from agricultural chemicals: A national perspective. Staff Report AGES870318. U.S. Dept. of Agriculture, Washington, DC.
- Peltier, W. H., and Weber, C. I. (Eds.) (1985). Methods for measuring acute toxicity of effluent to freshwater and marine organisms. EPA/600/4-85/013. U.S. Environmental Protection Agency, Cincinnati, OH.
- Richards, R. P., Kramer, J. W., Baker, D. B., and Krieger, K. A. (1987). Pesticides in rain water in the northeastern United States. *Nature (London)* 327, 129-131.
- Rowe, E. L., Ziobro, R. J., Wang, C. J. K., and Dence, C. W. (1982). The use of an alga *Chlorella pyrenoidesa* and a duckweed *Lemna perpusilla* as test organisms for toxicity bioassays of spent bleaching liquors and their compounds. *Environ. Pollut. (Ser. A)* 27, 289-296.
- Sahai, R., Roy, P. S., and Kaur, G. (1977). Comparative studies of toxicity of synthetic detergents on two common duckweeds at Gorakhpur. *Indian J. Ecol.* 4, 242-245.
- Said, M. J. M., Culley, D. D., Standifer, L. C., Epps, E. A., Myers, R. W., and Boney, S. A. (1979). Effect of harvest rate, waste loading, and stocking density on the yield of duckweeds. *Proc. World Maricult. Soc.* 10, 769-780.
- † Schindler, D. W. (1988). Effects of acid rain on freshwater ecosystems. *Science* 239, 149-157.
- Seto, M., Takahashi, Y., Ushijima, T., and Tazaki, T. (1979). Chlorotic death of *Lemna gibba* by cadmium in different concentrations of nutritional minerals. *Japan. J. Limnol.* 40, 61-65.
- Stanley, R. A., and Madewell, C. E. (1976). Chemical tolerance of *Lemna minor* L. Circular 2-72, TVA.

- Staves, R. P., and Knaus, R. M. (1975). Duckweed toxicity to aquatic plants. *Aquat. Bot.* 23, 261-273.
- Takemoto, B. K., and Noble, D. (1975). Carbon assimilation and frond growth of duckweed (*Lemna minor*) under low and high irradiance. *New Phytologist* 75, 101-108.
- Taraldsen, J. E., and Norberg-King, I. (1975). Duckweed (*Lemna minor*). *Environ. Contam. Toxicol.* 1, 1-14.
- Thomas, J. M., Skalske, J. P., Cline, S. A., Callahan, C. A., and Grover, R. P. (1985). Duckweed (*Lemna minor*) and its use in water quality assessment and determination of toxicants. *J. Fish. Aquat. Sci.* 40, 89-92.
- Van der Werff, M., and Pruyt, M. (1985). Duckweed (*Lemna minor*) as a bioindicator of water quality. *Chemosphere* 11, 727-739.
- Van Vierssen, W., and Prins, T. C. (1975). Duckweed (*Lemna minor*) as a bioindicator of water quality. *Chemosphere* 4, 101-108.
- Walbridge, L. T. (1977). A flow-through bioassay for duckweed (*Lemna minor*). EPA/600/3-77-108. U.S. Environmental Protection Agency, Washington, DC.
- Wang, W. (1986a). Toxicity tests of duckweed (*Lemna minor*). *Environ. Pollut. (Ser. B)* 11, 1-14.
- Wang, W. (1986b). The effect of river pollution on duckweed (*Lemna minor*). *Environ. Pollut. (Ser. B)* 11, 193-204.
- Wang, W. (1986c). Site-specific pollution assessment. U.S. Environmental Protection Agency, R810834. Illinois, U.S. Environmental Protection Agency, Springfield, IL.
- Wang, W. (1986d). Determination of toxicants in water using duckweed (*Lemna minor*). Report to Illinois DNR Survey Contract Report 405.
- Wang, W. (1987a). Chromate ion toxicity to duckweed (*Lemna minor*). *Toxicol. Chem.* 6, 409-414.
- Wang, W. (1987b). Toxicity of nickel to duckweed (*Lemna minor*). *Toxicol. Chem.* 6, 961-967.
- Wang, W. (1988). Site-specific bioassessment using duckweed (*Lemna minor*). EPA/600/3-88-122.
- Wang, W. Literature review on the use of duckweed (*Lemna minor*) as a bioindicator of water quality. Higher plants (common duckweed) submitted for publication.
- Wang, W., and Williams, J. M. (1985). Phytotoxicity tests. *Environ. Contam. Toxicol.* 1, 1-14.
- Wang, W., and Williams, J. M. (1986). Duckweed (*Lemna minor*) and millet for determining the toxicity of heavy metals in water. *Environ. Contam. Toxicol.* 1, 1-14.
- Wang, W. H., and Lay, J. P. (1985). Duckweed (*Lemna minor*) as a bioindicator of water quality. *Environ. Contam. Toxicol.* 1, 1-14.
- Wu, T. L., Lambert, L., Hasting, P., and Atrazine in the microsurface

- Bauer, R., Huber, W., and Sanz, M. (1985). Duckweed (*Lemna minor* L. Z. Pflanzenphysiologie) as a bioindicator of water quality. Study of 137 Cs adsorption on duckweed (*Lemna minor*). Birmingham, B. C., and Colmar, S. A. (1985). Duckweed (*Lemna minor*) as a bioindicator of water quality. *Environ. Contam. Toxicol.* 1, 1-14.
- Bowker, D. W., Duffield, A. N., and Callahan, C. A. (1985). Duckweed (*Lemna minor*) as a bioindicator of water quality. *Environ. Contam. Toxicol.* 1, 1-14.

- Staves, R. P., and Knaus, R. M. (1985). Chromium removal from water by three species of duckweeds. *Aquat. Bot.* 23, 261-273.
- Takemoto, B. K., and Noble, D. (1986). Differential sensitivity of duckweeds (*Lemnaceae*) to sulfite. I. Carbon assimilation and frond replication rate as factors influencing sulfite phytotoxicity under low and high irradiance. *New Phytol.* 103, 525-539.
- Taraldsen, J. E., and Norberg-King, T. A new method for determining effluent toxicity using duckweed (*Lemna minor*). *Environ. Toxicol. Chem.*, in press.
- Thomas, J. M., Skalske, J. P., Clire, J. F., McShane, M. C., Simpson, J. C., Miller, W. E., Peterson, S. A., Callahan, C. A., and Greene, J. C. (1986). Characterization of chemical waste site contamination and determination of its extent using bioassays. *Environ. Toxicol. Chem.* 5, 487-501.
- Threader, R. W., and Houston, A. H. (1983). Use of NaCl as a reference toxicant for goldfish. *Canad. J. Fish. Aquat. Sci.* 40, 89-92.
- Van der Werff, M., and Pruyt, M. J. (1982). Long-term effects of heavy metals on aquatic plants. *Chemosphere* 11, 727-739.
- Van Vierssen, W., and Prins, T. C. (1985). Relationship between the growth of algae and aquatic macrophytes in brackish waters. *Aquat. Bot.* 21, 165-179.
- Walbridge, L. T. (1977). A flow-through testing procedure with duckweed (*Lemna minor* L.). EPA-600/3-77-108. U.S. Environmental Protection Agency, Duluth, MN.
- Wang, W. (1986a). Toxicity tests of aquatic pollutants by using common duckweed. *Environ. Pollut. (Ser. B)* 11, 1-14.
- Wang, W. (1986b). The effect of river water on phytotoxicity of Ba, Cd, and Cr. *Environ. Pollut. (Ser. B)* 11, 193-204.
- Wang, W. (1986c). Site-specific phytotoxicity of heavy metals. Report to U.S. Environmental Protection Agency, R810834. Illinois State Water Survey Contract Report 403.
- Wang, W. (1986d). Determination of sediment toxicity in Waukegan Harbor through phytoassay methods. Report to Illinois Department of Energy and Natural Resources. Illinois State Water Survey Contract Report 405.
- Wang, W. (1987a). Chromate ion as a reference toxicant in aquatic phytotoxicity tests. *Environ. Toxicol. Chem.* 6, 409-414.
- Wang, W. (1987b). Toxicity of nickel to common duckweed (*Lemna minor*). *Environ. Toxicol. Chem.* 6, 961-967.
- Wang, W. (1988). Site-specific Ba toxicity to common duckweed, *Lemna minor*. *Aquat. Toxicol.* 12, 203-212.
- Wang, W. Literature review on the use of plants for toxicity assessment, submitted for publication.
- Wang, W. Higher plants (common duckweed, lettuce, rice) for effluent toxicity assessment, to be submitted for publication.
- Wang, W., and Williams, J. M. (1988). Screening and biomonitoring of industrial effluents using phytotoxicity tests. *Environ. Toxicol. Chem.* 7, 645-652.
- Wang, W., and Williams, J. M. (1990). The use of phytotoxicity tests (common duckweed, cabbage, and millet) for determining effluent toxicity. *Environ. Monit. Assess.*, 14, 59-69.
- Wang, W. H., and Lay, J. P. (1989). Fate and effects of salicylic acid compounds in freshwater system. *Ecotoxicol. Environ. Saf.* 17, 308-316.
- Wu, T. L., Lambert, L., Hasting, D., and Banning, D. (1980). Enrichment of the agricultural herbicide atrazine in the microsurface water of an estuary. *Bull. Environ. Contam. Toxicol.* 24, 411-412.

#### ADDITIONAL BIBLIOGRAPHY

- Bauer, R., Huber, W., and Sankhla, N. (1976). Effect of abscisic acid on photosynthesis in *Lemna minor* L. *Z. Pflanzenphysiol.* 77, 237-246.
- Bergamini, P. G., Palmas, G., Piantelli, F., Sani, M., Banditelli, P., Previtiera, M., and Sodi, F. (1979). Study of 137 Cs adsorption by *Lemna minor*. *Health Phys.* 37, 315-321.
- Birmingham, B. C., and Colman, B. (1983). Potential phytotoxicity of diquat accumulated by aquatic plants and sediments. *Water, Air, Soil Pollut.* 19, 123-131.
- Bowker, D. W., Duffield, A. N., and Denny, P. (1980). Methods for the isolation, sterilization and cultivation of Lemnaceae. *Freshwater Biol.* 10, 385-388.

- Clark, J. R., VanHassel, J. H., Nicholson, R. B., Cherry, D. S., and Cairns, J., Jr. (1981). Accumulation and depuration of metals by duckweed (*Lemna perpusilla*). *Ecotoxicol. Environ. Saf.* 5, 87-96.
- Coler, R. A. (undated). Statistical study of the duckweed Rhizosphere as an eco-assay tool. Water Resources Research Center. Univ. of Massachusetts Press, Amherst.
- DeLange, L. (1975). Gibbosity in the complex *Lemna gibba*/*Lemna minor*: Literature survey and ecological aspects. *Aq. Botan.* 1, 327-332.
- Ernst, W. H. D., and van der Werff, M. M. (1978). Aquatic angiosperms as indicators of copper pollution. *Arch. Hydrobiol.* 83, 356-366.
- Filbin, G. J., and Hough, R. A. (1979). The effects of excess copper sulfate on the metabolism of the duckweed *Lemna minor*. *Aquat. Bot.* 7, 79-86.
- Frankhauser, H., Brunhold, C., and Erisman, K. H. (1976). The influence of sublethal concentrations of sulfur dioxide on morphology, growth, and product yield of the duckweed *Lemna minor* L. *Oecologia* 23, 201-209.
- Haller, W. T., Sutton, D. L., and Barlowe, W. C. (1974). Effects of salinity on growth of several aquatic macrophytes. *Ecology* 55, 891-894.
- Kenaga, E., and Moolenaar, R. J. (1979). Fish and daphnia toxicity as surrogates for aquatic vascular plants and algae. *Environ. Sci. Technol.* 13, 1479-80.
- Laube, H. R., and Wohler, J. R. (1973). Studies on the decomposition of a duckweed (Lemnaceae) community. *Bull. Torrey Bot. Club* 100, 238-240.
- Lien, B. C., Walker, J. R. L., and Peters, J. A. (1980). The effect of fluoroacetate ("compound 1080") and fluoride upon duckweeds. *N. Zealand J. Sci.* 23, 179-183.
- Liu, L. C., and Maldonado, A. C. (1979). A bioassay method for detecting herbicide concentrations in water. *J. Agric. Univ. P.R.* 63, 80-83.
- Mestayer, C. R., Culley, D. D., Jr., Standifer, L. C., and Koonce, K. L. (1984). Solar energy conversion efficiency and growth aspects of the duckweed, *Spirodela punctata* (G. F. W. Mey.) Thompson. *Aquat. Bot.* 19, 157-170.
- Pandit, A. K., Kaul, V., and Fotedar, D. N. (1978). A preliminary study of duckweed ecology and its natural control. *Int. J. Ecol. Environ. Sci.* 4, 107-115.
- Rodgers, J. H., Cherry, D. S., and Guthrie, R. K. (1978). Cycling of elements in duckweed (*Lemna perpusilla*) in an ash settling basin and swamp drainage system. *Water Res.* 12, 765-770.
- Sirois, D. L. Evaluation of protocols for the assessment of phytotoxicity. ASTM STP 1091 Amer. Soc. Testing Materials, in press.
- Strother, S. (1980). Toxic effects of exogenous surbose on *Lemna minor* and some other angiosperms. *Ann. Bot. (London)* 47, 531-533.
- Strauss, R. (1976). The effects of different alkali salts on growth and mineral nutrient of *Lemna minor* L. *Int. Rev. Gesamten Hydrobiol.* 61, 673-676.
- Takemoto, B. K., Noble, R. D., and Harrington, H. M. (1986). Differential sensitivity of duckweeds (Lemnaceae) to sulphite. II. Thio production and hydrogen sulphide emission as factors influencing sulphite phytotoxicity under low and high irradiance. *New Phytol.* 103, 541-548.
- Tanaka, O., Nasu, Y., Yanase, D., Takimoto, A., and Kugimoto, M. (1982). pH dependence of the copper effect on flowering, growth, and chlorophyll content in *Lemna paucicostata* 6746. *Plant Cell Physiol.* 23, 1479-82.
- Walker, J. R. L., and Evans, S. (1978). Effects of quaternary ammonium compounds on some aquatic plants. *Marine Pollut. Bull.* 9, 136-137.
- Wang, W. (1984). Editorial: Uses of aquatic plants for ecotoxicology. *Environ. Int.* 10(3), i-iii.
- Wang, W. (1990). Characterization of phytotoxicity of metal engraving effluent samples. *Environ. Monit. Assess.*, 14, 45-50.

## Lung Can

F. J. TEWES,\* L.

\*INBIFO Institute of Biology  
 †Department of Community Medicine  
 Environmental Hygiene, Un.

To study the possible as and experimental materials: retrospective study of 2000 were interviewed concern to environmental pollutant significant increased lung brands of tea commonly c assay. Significantly eleva metabolic activation using that further research is ne ingestants taken at low de interactive effect between: cancer etiology. © 1990 A.

The risk to develop lu Yet recent toxicological for uptake and metaboli other means. For examp intramuscular, intraperit with subsequent toxic a compounds which have pulmonary neutrophils, epithelial glands. Exam; sensitivity, lung disease effects on the immune s Organic compounds g olised in the lung and pe (B(a)P) can be absorbed deposited in the lung (R In view of the above, ingested as well as airb formation is available o and tea.

Previous reports hav 1979; Tamura *et al.*, 19 found associations bet